

Spatial assignment of test sample

December 10, 2016

Contents

Input	1
Isotope values of test sample	1
Output	1
Model	1
Map of best fitted reference sample	2
Best fitted reference entries	3
Testing robustness of assignment: Wilcoxon signed rank test	3
P-values for the k nearest neighbours in Wilcoxon Test	4
Goodness of fit of test sample:	4

Input

Website Identifier: 11

Isotope values of test sample

Table 1: Isotope values of test sample


13C/12C	15N/14N	18O/16O	2H/1H	34S/32S
-27	9.7	17.5	-44.4	9.7

Output

Model

```
##
## Call:
## train.kknn(formula = fmla, data = ivory.train, kmax = 15, distance = 2, kernel = knl)
##
## Type of response variable: nominal
## Minimal misclassification: 0.3765867
## Best kernel: triangular
## Best k: 15
Classifier: country_code
```


Map of best fitted reference sample



Best fitted reference sample:

- Site: Dem. Rep. Congo
- Country: CD
- Region: Central Africa
- CITES: Appendix I
- Lat: -0.082275
- Lon: 14.573353

Assignment of test sample to nearest neighbours



Best fitted reference entries

Table 2: Details of best fitted reference entry (row 1) and other fitted entries within best classifier

lon	lat	location	13C/12C	15N/14N	18O/16O	2H/1H	34S/32S
14.57	-0.08	Dem. Rep. Congo	-24.9	10.3	16.4	-49.5	10.0
18.20	-1.55	Dem. Rep. Congo, Leopold ii meer	-25.9	10.9	17.1	-39.7	6.2
14.47	-0.01	Dem. Rep. Congo	-25.0	10.2	15.8	-49.9	10.1
23.55	-1.35	Dem. Rep. Congo, Moma	-25.6	10.8	16.9	-44.4	5.7
17.11	-1.10	Dem. Rep. Congo, Lukolela	-24.8	10.5	15.4	-43.4	8.4
21.45	-1.00	Dem. Rep. Congo, Itoko	-24.2	10.2	16.9	-42.9	6.8
13.02	-1.43	Dem. Rep. Congo, Itoko	-24.2	10.4	18.1	-38.4	7.0
23.20	-1.00	Dem. Rep. Congo, Tshuapa	-26.4	12.1	16.2	-46.3	5.7
27.37	1.40	Dem. Rep. Congo, Nepoko	-24.3	10.5	18.0	-34.9	7.4
29.51	1.32	Dem. Rep. Congo, Irumu	-24.5	10.1	16.3	-39.9	5.7
23.55	-1.35	Dem. Rep. Congo, Moma	-24.9	11.1	16.3	-48.5	5.6

Country of prediction: CD

Testing robustness of assignment: Wilcoxon signed rank test

If $p\text{-value} > 0.05$, the test concludes that the isotope signature of the test sample is similar to the respective nearest neighbour reference sample.

P-values for the k nearest neighbours in Wilcoxon Test

“0.000000490, 0.000000490, 0.000000103, 0.000000026, 0.000000026, 0.000000026, 0.000000026, 0.000000026,
0.000000026, 0.000000026, 0.000000026”

Goodness of fit of test sample:

- good fit: if $p > 0.05$ for at least two tested nearest neighbour reference samples;
- moderate fit: if $p > 0.05$ for at least one tested nearest neighbour reference samples;
- uncertain: if $p > 0.05$ for none of the tested nearest neighbour reference samples.

Assumption: At least two nearest reference samples are available.

Overall goodness of fit of test sample: “**uncertain**”