

# Spatial assignment of test sample

December 10, 2016

## Contents

<b>Input</b>	<b>1</b>
Isotope values of test sample . . . . .	1
<b>Output</b>	<b>1</b>
Model . . . . .	1
Map of best fitted reference sample . . . . .	2
Best fitted reference entries . . . . .	3
Testing robustness of assignment: Wilcoxon signed rank test . . . . .	4
P-values for the k nearest neighbours in Wilcoxon Test . . . . .	4
Goodness of fit of test sample: . . . . .	4

## Input

Website Identifier: 154

## Isotope values of test sample

Table 1: Isotope values of test sample

13C/12C	15N/14N	18O/16O	2H/1H	34S/32S
-26	12	17.8	-41.5	9.9


## Output

### Model

```
##  
## Call:  
## train.kknn(formula = fmla, data = ivory.train, kmax = 15, distance = 2, kernel = knl)  
##  
## Type of response variable: nominal  
## Minimal misclassification: 0.1889986  
## Best kernel: triangular  
## Best k: 14
```

Classifier: **region**


## Map of best fitted reference sample



### Best fitted reference sample:

- Site: Central African Republic
- Country: CF
- Region: Central Africa
- CITES: Appendix I
- Lat: 8
- Lon: 19

## Assignment of test sample to nearest neighbours



### Best fitted reference entries

Table 2: Details of best fitted reference entry (row 1) and other fitted entries within best classifier

lon	lat	location	13C/12C	15N/14N	18O/16O	2H/1H	34S/32S
19.00	8.00	Central African Republic	-25.7	12.8	19.8	-36.6	9.0
15.08	2.43	Cameroon, S dosten Kamerun N he Kika	-25.2	13.4	19.2	-36.1	9.1
18.20	-1.55	Dem. Rep. Congo, Leopold ii meer	-25.9	10.9	17.1	-39.7	6.2
19.00	8.00	Central African Republic	-25.6	12.8	19.7	-32.8	9.4
27.18	2.25	Dem. Rep. Congo, Medje	-24.2	11.9	17.4	-37.8	6.8
11.27	-3.78	Congo, Shot in Southern Congo, on the At	-24.7	11.6	15.8	-48.1	11.0
14.57	-0.08	Dem. Rep. Congo	-24.9	10.3	16.4	-49.5	10.0
23.55	-1.35	Dem. Rep. Congo, Moma	-25.6	10.8	16.9	-44.4	5.7
27.37	1.40	Dem. Rep. Congo, Nepoko	-24.3	10.5	18.0	-34.9	7.4
13.02	-1.43	Dem. Rep. Congo, Itoko	-24.2	10.4	18.1	-38.4	7.0
23.20	-1.00	Dem. Rep. Congo, Tshuapa	-26.4	12.1	16.2	-46.3	5.7
26.23	1.52	Dem. Rep. Congo, Panga	-24.6	12.1	17.1	-39.9	5.2
23.55	-1.35	Dem. Rep. Congo, Moma	-24.1	11.3	16.6	-40.9	6.3
17.11	-1.10	Dem. Rep. Congo, Lukolela	-24.8	10.5	15.4	-43.4	8.4

Region of prediction: Central Africa

## Testing robustness of assignment: Wilcoxon signed rank test

If p-value  $> 0.05$ , the test concludes that the isotope signature of the test sample is similar to the respective nearest neighbour reference sample.

### P-values for the k nearest neighbours in Wilcoxon Test

“0.6848475, 0.0055961, 0.0018749, 0.0015382, 0.0005277, 0.0005277, 0.0000019, 0.0000001, 0.0000001, 0.0000001, 0.0000001, 0.0000001, 0.0000001, 0.0000001”

### Goodness of fit of test sample:

- good fit: if  $p > 0.05$  for at least two tested nearest neighbour reference samples;
- moderate fit: if  $p > 0.05$  for at least one tested nearest neighbour reference samples;
- uncertain: if  $p > 0.05$  for none of the tested nearest neighbour reference samples.

Assumption: At least two nearest reference samples are available.

Overall goodness of fit of test sample: “**moderate fit**”